

Continuous glucose monitoring: Always Confirm glucose level with a fingerstick/meter prior to treatment

Hypoglycemia: Follow *Standards of Care for Diabetes Management in the School Setting – Colorado*, unless otherwise indicated here: _____

For severe symptoms: Administer Glucagon < 16 years old = 0.5 cc and > 16 years = 1.0 cc IM OR ____ mg(s) IM,

Disconnect Pump, Call 911

Hyperglycemia: Follow *Standards of Care for Diabetes Management in the School Setting – Colorado*, unless otherwise indicated here: _____

Ketone Testing per *Standards of Care for Diabetes Management in the School Setting – Colorado* OR Other: _____

Insulin Pump: Follow *Guidelines for Insulin Administration by School Staff, Diabetes Resource Nurses February 2013*

- Pump settings are established by the student's healthcare provider and should not be changed by the school staff. All setting changes to be made at home or by student providing self care as indicated on IHP.
- Internal safety features for the insulin pump should be active at all times while the student is at school - (Alarms set conservatively).

Insulin Pump Brand: _____ Type of Insulin in pump _____

Sensitivity/Correction Factor: _____ unit of insulin for every _____ mg/dl above the target blood glucose range

If blood glucose is less than _____ mg/dl, wait to give meal bolus until after meal

If blood glucose is greater than _____ mg/dl, deliver a correction bolus prior to eating

Insulin to Carbohydrate ratio _____ units of insulin per _____ grams of carbohydrate

Carbohydrate ratio for snack _____ units per _____ gm of carbs _____ am _____ pm

Bolus for carbohydrates should occur immediately Before lunch After lunch ½ bolus before & ½ bolus after

Other: _____

Parent/guardian authorized to increase or decrease insulin to carb ratio 1 unit +/- 5 grams of carbohydrates

Pump Malfunctions: Disconnect pump when malfunctioning

If pump is operational then the insulin dosing can be calculated by using the pump bolus calculator and then insulin given by injection

If pump is not operational:

Give insulin as indicated here:

Call Parent and Health Care Provider (for orders)